

# The zeta function of $\mathfrak{g}_{147A}$ counting ideals

## 1 Presentation

$\mathfrak{g}_{147A}$  has presentation

$$\left\langle x_1, x_2, x_3, x_4, x_5, x_6, x_7 \left| \begin{array}{l} [x_1, x_2] = x_4, [x_1, x_3] = x_5, [x_1, x_6] = x_7, \\ [x_2, x_5] = x_7, [x_3, x_4] = x_7 \end{array} \right. \right\rangle.$$

$\mathfrak{g}_{147A}$  has nilpotency class 3.

## 2 The local zeta function

The local zeta function was first calculated by Luke Woodward. It is

$$\zeta_{\mathfrak{g}_{147A},p}^{\triangleleft}(s) = \zeta_p(s)\zeta_p(s-1)\zeta_p(s-2)\zeta_p(s-3)\zeta_p(3s-4)\zeta_p(3s-5)\zeta_p(5s-8) \\ \times \zeta_p(7s-6)\zeta_p(6s-8)^{-1}.$$

$\zeta_{\mathfrak{g}_{147A}}^{\triangleleft}(s)$  is uniform.

## 3 Functional equation

The local zeta function satisfies the functional equation

$$\zeta_{\mathfrak{g}_{147A},p}^{\triangleleft}(s) \Big|_{p \rightarrow p^{-1}} = -p^{21-16s} \zeta_{\mathfrak{g}_{147A},p}^{\triangleleft}(s).$$

## 4 Abscissa of convergence and order of pole

The abscissa of convergence of  $\zeta_{\mathfrak{g}_{147A}}^{\triangleleft}(s)$  is 4, with a simple pole at  $s = 4$ .

## 5 Ghost zeta function

This zeta function is its own ghost.

## 6 Natural boundary

$\zeta_{\mathfrak{g}_{147A}}^{\triangleleft}(s)$  has meromorphic continuation to the whole of  $\mathbb{C}$ .

## 7 Notes

This ideal zeta function is identical to that of  $\mathfrak{g}_{147B}$ , though the Lie rings themselves are non-isomorphic.