

# The zeta function of $\mathfrak{g}_{6,4}$ counting ideals

## 1 Presentation

$\mathfrak{g}_{6,4}$  has presentation

$$\langle x_1, x_2, x_3, x_4, x_5, x_6 \mid [x_1, x_2] = x_5, [x_1, x_3] = x_6, [x_2, x_4] = x_6 \rangle.$$

$\mathfrak{g}_{6,4}$  has nilpotency class 2.

## 2 The local zeta function

The local zeta function was first calculated by Grunewald, Segal & Smith. It is

$$\zeta_{\mathfrak{g}_{6,4},p}^{\triangleleft}(s) = \zeta_p(s)\zeta_p(s-1)\zeta_p(s-2)\zeta_p(s-3)\zeta_p(3s-4)\zeta_p(5s-5)\zeta_p(6s-9) \\ \times \zeta_p(8s-9)^{-1}.$$

$\zeta_{\mathfrak{g}_{6,4}}^{\triangleleft}(s)$  is uniform.

## 3 Functional equation

The local zeta function satisfies the functional equation

$$\zeta_{\mathfrak{g}_{6,4},p}^{\triangleleft}(s) \Big|_{p \rightarrow p^{-1}} = p^{15-10s} \zeta_{\mathfrak{g}_{6,4},p}^{\triangleleft}(s).$$

## 4 Abscissa of convergence and order of pole

The abscissa of convergence of  $\zeta_{\mathfrak{g}_{6,4}}^{\triangleleft}(s)$  is 4, with a simple pole at  $s = 4$ .

## 5 Ghost zeta function

This zeta function is its own ghost.

## 6 Natural boundary

$\zeta_{\mathfrak{g}_{6,4}}^{\triangleleft}(s)$  has meromorphic continuation to the whole of  $\mathbb{C}$ .

## 7 Notes

The Lie ring is sometimes written as  $(F_{2,3}/\langle z \rangle) \cdot \mathbb{Z}$ .